

TABLE 6.2 SOME U.S. STATE STANDARDS FOR DISSOLVED OXYGEN

State	Surface water use classification				Comments	Reference ^a		
	Highest		Lowest					
	Daily average (mg/l)	Minimum (mg/l)	Daily average (mg/l)	Minimum (mg/l)				
Alabama	5	5	5	2	Applies @ 5 ft for depth > 10 ft and mid for 10 ft	1		
Georgia	6	5	5*	3	*Not less than value for at least $\frac{1}{24}$ hr	2		
Iowa	7*	5	4	4		3		
New Mexico	6	6	4	4		4		
Ohio	6	6	3	3	*During 4/1-6/15 and 9/16-12/31, not <6.5 mg/l as seasonal average	5		
Pennsylvania	7	3.5*	3	3	Diurnal decrease of 1 mg/l below standard allowed for no more than $\frac{1}{24}$ hr	6		
Texas	6	6	2	2		7		
Washington	9.5	4	4	4	Natural levels can be decreased by 0.2 mg/l	8		

^aReferences:

- (1) Alabama (1981).
- (2) Georgia (1980).
- (3) Iowa (1983).
- (4) New Mexico (1981).
- (5) Ohio (undated).
- (6) Pennsylvania (1979).
- (7) Texas (1981).
- (8) Washington (1982).

SOURCE : R.V. Thomann & T.A. Mueller
Harper Collins Publishers, 1987